New Jersey Department of Health Vaccine Preventable Disease Program

REQUEST FOR MEDICAL EXEMPTION FROM MANDATORY IMMUNIZATION

INSTRUCTIONS FOR COMPLETION

It is easiest to use the latest version of Adobe Reader DC. If you do not have the latest version, download and install the free software by visiting this webpage: https://get.adobe.com/reader/

- 1. Fill out the form completely. ALL form fields are required except where noted as being optional.
 - a. Enter the name of the Student and other identifying information.
 - b. Check off each vaccine for which an exemption is requested.
 - i. For each vaccine for which an exemption is requested, check to indicate whether the exemption is Temporary (indicate the date through which the exemption is valid) or Permanent.
 - ii. Check the ACIP contraindication/precaution applicable for each vaccine for which an exemption is requested.
 - c. If the contraindication/precaution is not included in Table 1, please put an "X" next to "Other" and fully explain. Please be sure that the contraindication/precaution does not appear in Table 2, that there is a valid contraindication/precaution noted for each vaccine for which an exemption is requested, and that the contraindication/precaution is consistent with ACIP/AAP guidelines and established national standards for vaccination practices.
- 2. Sign and date the Attestation Statement
- 3. Provide a copy to the person requesting the medical exemption or directly to the school, preschool or child care center.
- 4. Keep a copy of the form for your records.

Name of Student (first, middle, last)			Date of Birth
Name of Parent/Guardian (if under 18) (first, middle, last)			Primary Phone
Patient/Parent Home Address		Address Line 2	
City	State		Zip Code
Patient/Parent Email Address			

Medical contraindications and precautions for immunizations are based on the most recent General Recommendations of the Advisory Committee on Immunization Practices (ACIP), available at

https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html or https://redbook.solutions.aap.org/redbook.aspx

Please check the website to ensure that you are reviewing the most recent ACIP information. Please note that the presence of a moderate to severe acute illness with or without fever is a precaution to administration of all vaccines However, as acute illnesses are short-lived, medical exemptions should not be submitted for this indication.

IMM-53 AUG 25

Table 1. A		tions and Precautions to Vaccination for Mandatory Vaccines
Vaccine	Exemption Length	ACIP Contraindications and Precautions (CHECK ALL THAT APPLY)
DTaP, Tdap	Temporary through:	Contraindications
	Permanent	Severe Allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
		Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures) not attributable to another identifiable cause within 7 days of administration of a previous dose of DTP, DTaP, or Tdap
		Precautions
		Progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy; defer DTaP or Tdap until neurologic status clarified and stabilized
		Guillain-Barré syndrome < 6 weeks after previous dose of tetanus-toxoid- containing vaccine
		History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus toxoid-containing vaccine
DT, Td	Temporary through:	Contraindications
	Permanent	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
		Precautions
		Guillain-Barré syndrome < 6 weeks after a previous dose of tetanus- toxoid-containing vaccine.
		History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria- or tetanus toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid-containing vaccine
Haemophilus influenzae type b (Hib)	Temporary through:	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a
, ,	Permanent	vaccine component

Vaccine	Exemption Length	ACIP Contraindications and Precautions (CHECK ALL THAT APPLY)
Hepatitis B (HepB)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Hypersensitivity to yeast
Inactivated poliovirus vaccine (IPV)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Precautions Pregnancy
Influenza, inactivated injectable (IIV)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Precautions Guillain-Barré syndrome < 6 weeks after a previous dose of tetanus- toxoid-containing vaccine. Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, or recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting, under the supervision of a healthcare provider who is able to recognize and manage severe allergic conditions)
Influenza recombinant (RIV)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Precautions Guillian-Barré syndrome < 6 weeks after a previous dose of influenza vaccine

Vaccine	Exemption Length	ACIP Contraindications and Precautions (CHECK ALL THAT APPLY)
MMR	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Pregnancy Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with human immunodeficiency virus [HIV] infection who are severely immunocompromised) Family history of congenital or hereditary immunodeficiency in first-degree relatives (e.g., parents and siblings), unless the immune competence of the potential vaccine recipient has been substantiated clinically or verified by a laboratory test Precautions Recent (≤ 11 months) receipt of antibody-containing blood product (specific interval depends on product) History of thrombocytopenia or thrombocytopenic purpura Need for tuberculin skin testing or interferon gamma release assay (IGRA) testing
Meningococcal (MenACWY)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
Meningococcal (MenB)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Precautions Pregnancy

Vaccine	Exemption Length	ACIP Contraindications and Precautions (CHECK ALL THAT APPLY)
Pneumococcal (PCV13)	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid-containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid-containing vaccine), including yeast
Varicella	Temporary through: Permanent	Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or persons with HIV infection who are severely immunocompromised Pregnancy Family history of congenital or hereditary immunodeficency in first-degree relatives (e.g., parents and siblings), unless the immune competence of the potential vaccine recipient has been substantiated clinically or verified by a laboratory test Precautions Recent (≤ 11 months) receipt of antibody-containing blood product (specific interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination; avoid use of these antiviral drugs for 14 days after vaccination) Use of aspirin or aspirin-containing products

Other: Please explain fully and attach additional sheets as necessary. Please be sure to check Table 2 below to ensure that the condition is not one incorrectly perceived as a contraindication or precaution.

Attestation				
am a physician (M.D. or D.O.) licenses to practice medicine in a jurisdiction of the United States or an advanced practice nurse licensed in a jurisdiction of the United States.				
affirm that the stated contraine established national standard supporting medical document	dicatio s for v ation.	n(s)/precaution(s) is enul accination practices. I un I also understand that an	merated b derstand t y misrepre	indications and Precautions and y the ACIP and consistent with hat I might be required to submit esentation might result in referral te licensing/regulatory agency.
Healthcare Provider Name (please print) Specialty				
NPI Number		License Number	·	State of Licensure
Phone	Fax		Email	
Address				
City		State		Zip Code

Date

Signature

	of Conditions incorrectly perceived as contraindications or precuations to vaccination* nes may be given under these conditions)
Vaccine	Conditions incorrectly perceived as contraindications and precautions to vaccines (i.e., vaccines may be given under these conditions)
General for MMR, Hib, HepB, Varicella, PCV13, MenACWY	 History of Guillain-Barré syndrome Recent exposure to an infectious disease History of penicillin allergy, other nonvaccine allergies, relatives with allergies, or receiving allergen extract immunotherapy
DTaP	 Fever within 48 hours after vaccination with a previous dose of DTP or DTaP Collapse or shock like state (i.e., hypotonic hyporesponsive episode) within 48 hours after receiving a previous dose of DTP/DTaP Seizure ≤ 3 days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting ≥ 3 hours within 48 hours after receiving a previous dose of DTP/DTaP Family history of seizures Family history of sudden infant death syndrome Family history of an adverse event after DTP/DTaP Stable neurologic conditions (e.g., cerebral palsy, well-controlled seizures, or developmental delay)
Hepatitis B (HepB)	 Pregnancy Autoimmune disease (e.g., systemic lupus erythematosus or rheumatoid arthritis)
Influenza, inactivated injectable (IIV)	Nonsevere (e.g., contact) allergy to latex, thimerosal, or egg
MMR	 Breastfeeding Pregnancy of recipient's mother or other close or household contact Recipient is female of child-bearing age Immunodeficient family member or household contact Asymptomatic or mildly symptomatic HIV infection Allergy to eggs
Tdap	 History of fever of ≥ 40.5° C (≥ 105° F) for < 48 hours after vaccination with previous dose of DTP/DTaP History of collapse or shock-like state (hypotonic hyporesponsive episode) within 48 hours after receiving a previous dose of DTP/DTaP History of persistent, inconsolable crying lasting > 3 hours within 48 hours of receiving a previous dose of DTP/DTaP History of extensive limb swelling after DTP/DTaP/Td that is not an Arthus-type reaction History of stable neurologic disorder Immunosuppression
Varicella	Pregnancy of recipient's mother or other close or household contact Immunodeficient family member or household contact Asymptomatic or mildly symptomatic HIV infection

^{*} For a complete list of conditions, please review the ACIP Guide to Contraindications and Precautions accessible at https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html

Humoral immunodeficiency (e.g., agammaglobulinemia)